

## Hiroshima survivors exposed to very low doses of A-bomb primary radiation showed a high risk for cancers

Masaru Miyao · Tomoyuki Watanabe ·  
Ryumon Honda · Yuichi Yamada

Published online: 4 April 2009  
© The Japanese Society for Hygiene 2009

**Erratum to: Environ Health Prev Med (2008) 13:264–270**  
**DOI 10.1007/s12199-008-0039-8**

Our paper contained the following two errors. First, on pages 266 and 267, the phrase “person-years” should be “person” in the Table 1 and 2 footnotes “<sup>a</sup> Observed number of deaths (person-years)” and “<sup>b</sup> Expected number

of deaths (person-years)”. Second, on page 267, the 95% confidential intervals (95% CI) for the SMR-O in the female very low dose category (<0.005 Sv) due to leukemia was “(0.469–1.945)” in the Table 2. However, we should have been listed as “(0.503–2.084)”. We regret any confusion these errors might have caused.

---

The online version of the original article can be found under doi:[10.1007/s12199-008-0039-8](https://doi.org/10.1007/s12199-008-0039-8).

---

M. Miyao (✉)  
Graduate School of Information Science, Nagoya University,  
Furo-cho, Chikusa-ku, Nagoya 464-8601, Japan  
e-mail: [mmiyao@med.nagoya-u.ac.jp](mailto:mmiyao@med.nagoya-u.ac.jp)

T. Watanabe  
Department of Health and Nutrition,  
Faculty of Psychological and Physical Science,  
Aichi Gakuin University, Nisshin, Japan

R. Honda · Y. Yamada  
Department of Social and Environmental Medicine (Hygiene),  
Kanazawa Medical University, Ishikawa, Japan