

CORRECTION

Open Access



Correction to: Association between *NAT2*, *CYP1A1*, and *CYP1A2* genotypes, heterocyclic aromatic amines, and prostate cancer risk: a case control study in Japan

Masahide Koda^{1*}, Motoki Iwasaki², Yuko Yamano³, Xi Lu¹ and Takahiko Katoh¹

Correction

Following publication of the original article [1], the authors reported a correction in the units in the methods section under “Subjects”

Originally, the sentence was stated as:

(ii) exhibited no evidence of prostate cancer as determined by blood test (PSA < 4 mg/dL)

The correct sentence is:

(ii) exhibited no evidence of prostate cancer as determined by blood test (PSA < 4 ng/ml)

Author details

¹Department of Public Health, Faculty of Life Sciences, Kumamoto University, 1-1-1 Honjou, Chuo-ku, Kumamoto 860-8556, Japan. ²Division of Epidemiology, Center for Public Health Sciences, National Cancer Center, Tokyo, Japan. ³Department of Hygiene and Preventive Medicine, Showa University School of Medicine, Tokyo, Japan.

Published online: 04 July 2018

Reference

1. Koda M, et al. Association between *NAT2*, *CYP1A1*, and *CYP1A2* genotypes, heterocyclic aromatic amines, and prostate cancer risk: a case control study in Japan. *Environ Health Prev Med*. 2017;22:72. <https://doi.org/10.1186/s12199-017-0681-0>.

* Correspondence: shoei05-kmi@umin.ac.jp

¹Department of Public Health, Faculty of Life Sciences, Kumamoto University, 1-1-1 Honjou, Chuo-ku, Kumamoto 860-8556, Japan

